

PRESCRIBING INFORMATION

(Please refer to the full Summary of Product Characteristics for detailed information)

balance 1.5% glucose 1.25 mmol/l calcium, solution for peritoneal dialysis MA No: PL 13689/0011
balance 2.3% glucose 1.25 mmol/l calcium, solution for peritoneal dialysis MA No: PL 13689/0013
balance 4.25% glucose 1.25 mmol/l calcium, solution for peritoneal dialysis MA No: PL 13689/0012

Composition:

1 litre of the neutral ready-to-use solution contains:

Active substances	balance 1.5% glucose, 1.25 mmol/l calcium	balance 2.3% glucose, 1.25 mmol/l calcium	balance 4.25% glucose, 1.25 mmol/l calcium
Sodium chloride	5.640	5.640	5.640
Sodium (S) lactate solution (sodium (s) lactate)	7.85g (3.925g)	7.85g (3.925g)	7.85g (3.925g)
Calcium chloride dihydrate	0.1838	0.1838	0.1838
Magnesium chloride hexahydrate	0.1017	0.1017	0.1017
Glucose monohydrate (anhydrous glucose)	16.5g (15.0g)	25.0g (22.73g)	46.75g (42.5g)

1 litre of the neutral ready-to-use solution contains:

Active substances in mmol	balance 1.5% glucose, 1.25 mmol/l calcium	balance 2.3% glucose, 1.25 mmol/l calcium	balance 4.25% glucose, 1.25 mmol/l calcium
Na ⁺	134	134	134
Ca ²⁺	1.25	1.25	1.25
Mg ²⁺	0.5	0.5	0.5
Cl ⁻	100.5	100.5	100.5
(s)-Lactate	35	35	35
Glucose	83.2	126.1	235.8
Theoretical osmolarity (mosm/l)	356	399	509
pH	7.0	7.0	7.0

Indications: End-stage (decompensated) chronic renal failure of any origin which can be treated with peritoneal dialysis.

Dosage and Administration: The mode of therapy, frequency of administration, and dwell time required will be specified by the attending physician. **Continuous ambulatory peritoneal dialysis (CAPD): Adults:** Unless otherwise prescribed, patients will receive an infusion of 2000 ml solution per exchange 4 times a day. After a dwell time between 2 and 10 hours the solution will be drained. Adjustment of dosage, volume and number of exchanges will be necessary for individual patients. **Paediatric population:** In children the solution volume per exchange should be prescribed according to age and body surface area (BSA). For initial prescription, the volume per exchange should be 600-800ml/m² BSA with 4 (sometimes 3 or 5) exchanges per day. It can be increased up to 1000-1200ml/m² BSA depending on tolerance, age and residual renal function. **Automated peritoneal dialysis (APD)** a machine (cycler) is used for intermittent or continuous cyclic peritoneal dialysis, the use of larger bags is recommended providing more than one solution exchange. **Adults:** Typically, patients spend 8-10 hours a night cycling. Dwell volumes range from 1500 to 3000 ml and the number of cycles usually varies from 3 to 10 per night. The amount of fluid used is typically between 10 and 18 litres but can range from 6 to 30 litres. The cylinder therapy at night is usually combined with 1 or 2 exchanges during the daytime. **Paediatric population:** The volume per exchange should be 800-1000 ml/m² BSA with 5-10 exchanges overnight. It can be increased up to 1400 ml/m² BSA depending on tolerance, age and residual renal function. **Elderly population:** There are no special dosage recommendations for the elderly.

Method of administration: Before performing peritoneal dialysis at home, the patient must be trained appropriately, must practice the technique and be shown to be proficient. The training should be performed by qualified personnel. The attending physician must ensure that the patient masters the handling techniques sufficiently before being discharged to carry out peritoneal dialysis at home. In case of any problems or uncertainty the attending physician should be contacted. **Continuous ambulatory peritoneal dialysis (CAPD): stay-safe bag:** The solution bag is first warmed up to body temperature. The appropriate dose is infused in the peritoneal cavity using a peritoneal catheter over 5 - 20 minutes. Depending on physician's instructions, the dose should dwell in the peritoneal cavity for 2 to 10 hours (equilibrium time), and then be drained. **Automated peritoneal dialysis (APD): sleep-safe bag:** The connectors of the prescribed sleep-safe solution bags are inserted in the free tray ports and then automatically connected to the tubing set by the cycler. The cycler checks the bar codes of the solution bags and gives an alarm when the bags do not comply with the prescription stored in the cycler. After this check the tubing set can be connected to the patient's catheter extension and the treatment be started. The sleep-safe solution is automatically warmed up to body temperature by the cycler during the inflow into the abdominal cavity. **Contraindications:** For this specific peritoneal dialysis solution **balance 1.5%/2.3% glucose, 1.25 mmol/l calcium** must not be used in patients with lactic acidosis, severe hypokalaemia and severe hypocalcaemia. **balance 4.25% glucose, 1.25 mmol/l calcium** must not be used in patients with lactic acidosis, severe hypokalaemia, severe hypocalcaemia, hypovolaemia and arterial hypotension. **For peritoneal dialysis in general -** recent abdominal surgery or injury, extensive inflammatory conditions of the abdominal skin, inflammatory bowel diseases, peritonitis, internal or external abdominal fistula, umbilical, inguinal or other abdominal hernia, intra-abdominal tumours, ileus, pulmonary disease, sepsis, extreme hyperlipidaemia, in rare cases of uraemia, cachexia and severe weight loss.

Contraindications: Peritoneal dialysis should not be commenced if any of the following are present: recent abdominal surgery or injury, extensive inflammatory conditions of the abdominal skin, inflammatory bowel diseases, peritonitis, internal or external abdominal fistula, umbilical, inguinal or other abdominal hernia & intra-abdominal tumours.

Special warnings and precautions for use: Encapsulating peritoneal sclerosis is considered to be a known, rare complication of peritoneal dialysis therapy which can infrequently lead to fatal outcome. The solution for peritoneal dialysis must not be used for intravenous infusion. This solution may only be administered after careful benefit-risk assessment in: **balance 1.5%/2.3%/4.25% glucose, 1.25 mmol/l calcium - patients with hyperparathyroidism:** Therapy should include the administration of calcium-containing phosphate binders and/or vitamin D to ensure adequate enteral calcium supply. **Hypocalcaemia:** It may be necessary to use a peritoneal dialysis solution with a higher calcium concentration either temporarily or permanently, in case an adequate enteral supply of calcium, by calcium-containing phosphate binders and/or vitamin D, is not possible. **balance 1.5%/2.3% glucose, 1.25 mmol/l calcium and 4.25% glucose, 1.25 mmol/l calcium - loss of electrolytes due to vomiting and/or diarrhoea** (a temporary change to a peritoneal dialysis solution containing potassium might then become necessary). **Patients receiving digitalis therapy:** Regular monitoring of the serum potassium level is mandatory. Severe hypokalaemia may necessitate the use of a potassium-containing dialysis solution together with dietary counselling. **Patients with large polycystic kidneys:** a loss of proteins, amino acids, and water-soluble vitamins occurs during peritoneal dialysis. To avoid deficiencies an adequate diet or supplementation should be ensured. **Elderly Population -** The increased incidence of hernia should be considered in the elderly prior to the start of peritoneal dialysis. **Interactions:** The use of this peritoneal dialysis solution can lead to a loss of efficacy of other medicinal products if these are dialysable through the peritoneal membrane.

Undesirable effects:

Peritoneal dialysis solution: Endocrine disorders - balance 1.5%/2.3%/4.25% glucose, 1.25 mmol/l calcium - Secondary hyperparathyroidism with potential disturbances of the bone metabolism (not known). **Metabolism and nutrition disorders -** Increased blood sugar levels (common ≥1/100 to <1/10), Hyperlipidaemia (common), Increase in body weight due to the continuous uptake of glucose from the peritoneal dialysis solution (common). **Cardiac disorders -** Tachycardia (uncommon ≥1/1000 to <1/100), Hypotension (uncommon). Hypertension (uncommon). **Respiratory, thoracic and mediastinal disorders -** Dyspnoea (uncommon). **Renal and urinary disorders - balance 1.5%/2.3%/4.25% glucose, 1.25 mmol/l calcium -** Electrolyte disturbances (very common ≥1/10), Hypocalcaemia (uncommon)

Treatment mode: Infections and infestations - Peritonitis (very common), Skin exit site and tunnel infections (very common) and Sepsis (very rare <1/10,000). **Respiratory, thoracic and mediastinal disorders -** Shoulder pain (common) and Dyspnoea caused by the elevated diaphragm (not known). **Gastrointestinal disorders -** Hernia (very common), Abdominal distension and sensation of fullness (common), Diarrhoea (uncommon), Constipation (uncommon) and Encapsulating peritoneal sclerosis (not known). **General disorders and administration/catheter site conditions -** Redness, oedema, exudations, crusts and pain at the catheter exit site (very common), In- and outflow disturbances of the dialysis solution (common), General malaise (not known). **Presentations: stay-safe:** The stay-safe system is provided as a double-bag system consisting of the double-chamber solution bag. **sleep-safe:** The sleep-safe system is provided as a single-bag system consisting of the double-chamber solution bag.

(Please refer to the full Summary of Product Characteristics for detailed information)

Legal Category: POM

Marketing Authorisation Holder: Fresenius Medical Care Deutschland GmbH D-61346 Bad Homburg Germany.

Presentation and NHS cost (excl. VAT): stay.safe balance 2000ml bag £7.50, 2500ml bag £8.30, sleep.safe balance 5000ml bag £11.60

Date of revision: February 2026



Fresenius Medical Care (UK) Ltd (local supplier)
Nunn Brook Road,
Huthwaite, Sutton-In-Ashfield
Nottinghamshire, NG17 2HU
Telephone: 0800 001 4499

Adverse events should be reported. Reporting forms and information can be found via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard. Adverse events should also be reported to Fresenius Medical Care Vigilance team on 01623 445 215